

# Evaluation of Tillering Behavior and Yielding Ability of Different Rice Varieties under Unpuddled Conditions

**K. Srimathi <sup>a</sup> and E. Subramanian <sup>b</sup>**

<sup>a</sup> Department of Agronomy, TNAU, Coimbatore – 641 003, India.

<sup>b</sup> Department of Agronomy, AC & RI, Killikulam – 641 003, India.

## **Authors' contributions**

*This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.*

## **Article Information**

DOI: 10.9734/CJAST/2022/v41i2831792

## **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89853>

**Original Research Article**

**Received 28 May 2022**  
**Accepted 01 August 2022**  
**Published 08 August 2022**

## **ABSTRACT**

A field trial was conducted at Central Farm of Agricultural College and Research Institute, Madurai during *Kharif*, 2017 on sandy clay loam soils to evaluate the effect of various sowing windows on tillering performance and yielding ability of three different varieties under unpuddled condition. The study was evaluated in a Split plot design with three replications and a treatment consisting of two dates of sowing (July first and second fortnight) assigned in main plots and three varieties *viz.*, ADT 48, MDU 5 and CO 51 in two methods of planting (Direct seeded rice (DSR) and Unpuddled Transplanted rice (UPTR)) in sub plots. The treatments were compared with conventional planting of short duration variety (CO 51) under late release of water from July first and second fortnight (absolute control 1 & 2). The results revealed that, tiller production was not significantly influenced by sowing dates but effective tillers and yield was more in early sowing than late sowing. Among the method of planting with different varieties, direct sowing of CO 51 produced more tillers like primary, secondary tillers  $m^{-2}$  at various growth stages. The tertiary tiller production was higher in the UPTR method than direct sowing. As a result of more number of panicles, higher yield was achieved in direct sowing of CO 51 which produced a grain yield of  $5987 \text{ kg ha}^{-1}$  and straw yield of  $8130 \text{ kg ha}^{-1}$  over early and very early duration varieties *viz.*,

<sup>o</sup> Ph.D Scholar;

<sup>#</sup> Assistant Professor;

<sup>\*</sup>Corresponding author: E-mail: [srimathiraj95@gmail.com](mailto:srimathiraj95@gmail.com);

MDU 5 and ADT 48 under both method of planting. Therefore, CO 51 was proved its supremacy to produce more tillers and give higher yield under direct sowing method than unpuddled condition transplanting.

**Keywords:** Direct sowing; UPTR; varieties; sowing window; tillering performance.

## 1. INTRODUCTION

Rice (*Oryza sativa* L.) is a prime member of graminiae family and predominant survival food crop for majority of the world population. Rice cultivation and its productivity is being a challenge in the upcoming decades due to potential climate change. Recently, rice production in India is limited due to water scarcity and monsoon failure [1]. Water is becoming increasingly scarce in the traditional river valley command area because of many reasons like erratic monsoon behavior, deteriorating hydrological balance and increasing demand for non-agricultural purpose [2].

The traditional practice is raising short duration variety in the first season followed by medium duration variety in the second season, and there are benefited by South West Monsoon and North East Monsoon rains respectively. Changes in on set of South West Monsoon and inadequate level of water in the dam, makes cropping difficult. In some years, the water released for first crop is delayed even up to August consequently, the transplanting of first crop get delayed which results in delayed transplanting of second crop (medium duration). This situation makes the second rice crop to suffer for the want of sufficient water chiefly at flowering and grain filling stages. This shows that some of the *Kharif* area (double crop) has been shifted to *rabi* (single crop) mainly consequent of late withdrawal of water from the reservoir. If this situation continues, it is feared that *Kharif* crop would completely vanish.

Direct sowing of rice seeds in unpuddled land offers an useful option to reduce the limitations of transplanted paddy and also offers one such water saving, less labour usage technology. This concept is mainly suited for irrigated lowlands, where supplemental irrigation is available for crop cultivation [3]. Likewise the Unpuddled Transplanted Rice (UPTR) is gaining momentum, wherein 30 per cent of water, which is utilized for field preparation, has been reduced and the turnaround period between crops also get reduced. However at the same time, availability of short, early and very early duration, photo-

insensitive, semi-dwarf and high yielding rice varieties are a boon for direct seeded technique. Direct seeded rice matures seven to ten days earlier than transplanted rice, which gives more importance in intensive cropping [4]. In this context, very early, early and short duration rice varieties viz., ADT 48, MDU 5 and CO 51 have the duration of 95, 100-105 and 110 days respectively may suit for the delayed sowing especially at first season due to late receipt of canal water. Further, it makes the second season rice without any delay in yield penalty. Under these circumstances, the study was carried out to evaluate the promising variety for late receipt of canal water in *Kharif* season to enhance the rice productivity.

## 2. MATERIALS AND METHODS

A field experiment was carried out in *Kharif*, 2017 at the Agricultural College and Research Institute, Madurai, Tamil Nadu situated at 9°54' N latitude and 78°54' E longitude with an altitude of 147 m above mean sea level. The track is located in the southern agro climatic zone of Tamil Nadu. The soil of the experiment field was sandy clay loam in texture. The trial was laid out in a split plot design and replicated thrice. The treatments combinations comprised two dates of sowing, viz., first and second fortnight of July in main plots and two methods of planting with three varieties (Direct sowing and Unpuddled Transplanted rice (UPTR) with CO 51, MDU 5 and ADT 48 rice varieties) in sub plots. The seeds were soaked in water for 12 hours and incubated for 10 hours. Sprouted seeds were line sown with a spacing of 20 x10 cm and (seedlings were already raised in community nursery) eighteen days old seedlings were transplanted with same spacing in a saturated soil on the same day (July first and second fortnight). Irrigation was provided immediately after sowing to hasten the germination and establishment. Subsequent irrigations were given to maintain moist condition and need based plant protection was given. The tillers production recorded for each treatment by counting the uprooted plant tillers at active tillering stage, panicle initiation stage, flowering stage and at harvest. Primary, secondary and tertiary tillers hill<sup>-1</sup> were also

recorded. The yield attributes and yield were recorded and statistically analyzed at 5 % level of significance [5].

### 3. RESULTS AND DISCUSSION

#### 3.1 Tiller Production $m^{-2}$

The data on primary, secondary, tertiary and total tillers per unit area recorded at different growth stages are presented in Tables 1 and 2. The trend in variation of tiller production per unit area was not significantly differed with respect to date of sowing only. Method of planting with different varieties produced significant variation at all growth stages of crop. The number of tillers  $m^{-2}$  is one of the most important growth parameters which influence the grain yield directly. It is apparent from the data indicated that tiller production  $m^{-2}$  was increased very fast up to panicle initiation stage, thereafter it became very slow and follow the Mitscherlich concept of increases with decreasing trend.

With regard to the method of planting with different varieties, significant variation was noticed in primary, secondary, tertiary and total tillers  $m^{-2}$ . During the active tillering stage, direct sowing of CO 51 resulted in significantly higher primary (283.9) and secondary tillers  $m^{-2}$  (199.4) and this was on par with direct sowing of MDU 5

and UPTR of CO 51 rice variety. The significantly lower numbers of primary and secondary tillers  $m^{-2}$  (199.2 and 148.1) was recorded in case of ADT 48 under UPTR. At panicle initiation stage, primary tiller production was more in direct sowing than the unpuddled transplanting. Direct sowing of CO 51 produced more primary tillers (367.9) and lower number of primary tillers (247.4) was observed under UPTR of ADT 48, which was on par with UPTR of MDU 5. With respect to secondary tillers  $m^{-2}$  was more in CO 51 under the direct sowing (375.2) and less in ADT 48 under direct sowing and UPTR method (310.3 and 297.2, respectively). Transplanted rice produced more tertiary tillers than direct sowing method. A higher tertiary tillers  $m^{-2}$  (75.6) was recorded in UPTR of CO 51 and direct sowing of ADT 48 produced significantly lower number of tertiary tillers  $m^{-2}$  (46.7).

During the flowering stage, direct sowing of CO 51 produced more primary tillers  $m^{-2}$  (364.6). The lower number of primary tillers was recorded in ADT 48 under direct sowing and UPTR. A higher number of secondary tillers  $m^{-2}$  was noticed under direct sowing of CO 51 (441.2). The lowest number of secondary tillers (347.4) was observed under ADT 48 in UPTR. With respect to tertiary tillers  $m^{-2}$ , the rice variety CO 51 produced more tertiary tillers  $m^{-2}$  followed by MDU 5.

**Table 1. Date of sowing and method of planting on primary, secondary, tertiary and total tillers  $m^{-2}$  at AT and PI stages of rice**

Treatment	Active tillering stage			Panicle initiation stage			
	Primary	Secondary	Total	Primary	Secondary	Tertiary	Total
<b>Main plot : Date of sowing</b>							
D <sub>1</sub>	243.7	174.3	418.1	311.8	348.8	63.3	725.6
D <sub>2</sub>	231.8	166.8	399.5	289.3	310.9	54.3	655.0
<b>Mean</b>	237.8	170.5	408.8	300.5	329.9	58.8	690.3
<b>SEd</b>	<b>7.86</b>	<b>3.18</b>	<b>7.11</b>	<b>5.29</b>	<b>10.1</b>	<b>2.76</b>	<b>17.3</b>
<b>CD (p=0.05)</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>
<b>Sub plot : Method of planting/sowing with different varieties</b>							
T <sub>1</sub>	283.9	199.4	483.2	367.9	375.2	65.5	808.6
T <sub>2</sub>	242.2	171.0	416.6	316.7	317.1	51.8	685.5
T <sub>3</sub>	220.6	159.5	375.8	290.6	310.3	46.7	647.6
T <sub>4</sub>	251.7	178.3	431.0	315.7	357.7	75.6	749.0
T <sub>5</sub>	228.8	166.9	398.9	264.9	321.6	59.1	653.9
T <sub>6</sub>	199.2	148.1	347.3	247.4	297.2	50.8	597.0
<b>Mean</b>	237.8	170.5	408.8	300.5	329.9	58.2	690.3
<b>SEd</b>	<b>7.04</b>	<b>6.67</b>	<b>11.5</b>	<b>10.5</b>	<b>11.6</b>	<b>4.54</b>	<b>17.9</b>
<b>CD (p=0.05)</b>	<b>14.2</b>	<b>13.4</b>	<b>23.2</b>	<b>21.1</b>	<b>23.3</b>	<b>9.16</b>	<b>36.2</b>

\*AT – Active Tillering stage; \*PI – Panicle Initiation stage

**Table 2. Date of sowing and method of planting on primary, secondary, tertiary and total tillers m<sup>2</sup> at flowering and harvest stages of rice**


Treatment	Flowering stage				Harvest stage			
	Primary	Secondary	Tertiary	Total	Primary	Secondary	Tertiary	Total
<b>Main plot : Date of sowing</b>								
D <sub>1</sub>	323.4	390.8	94.8	813.6	335.3	412.0	114.8	879.7
D <sub>2</sub>	309.7	381.1	89.0	778.3	322.1	391.4	99.6	797.8
<b>Mean</b>	316.6	386.0	91.9	795.9	328.6	402.0	107.2	838.7
<b>SEd</b>	<b>6.56</b>	<b>15.8</b>	<b>2.71</b>	<b>14.9</b>	<b>7.23</b>	<b>6.99</b>	<b>5.00</b>	<b>17.1</b>
<b>CD (p=0.05)</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>	<b>NS</b>
<b>Sub plot : Method of planting/sowing with different varieties</b>								
T <sub>1</sub>	364.6	441.2	113.9	927.7	384.9	458.7	122.5	966.2
T <sub>2</sub>	313.4	390.8	90.6	798.1	321.8	394.6	101.1	804.9
T <sub>3</sub>	302.8	364.3	87.5	737.8	305.4	379.1	90.6	787.3
T <sub>4</sub>	324.8	397.1	93.8	824.6	344.7	428.6	125.5	890.3
T <sub>5</sub>	312.1	375.1	87.8	781.6	319.6	384.6	109.9	834.5
T <sub>6</sub>	281.7	347.4	77.9	705.8	296.9	366.5	96.7	749.4
<b>Mean</b>	316.6	386.0	91.9	795.9	328.6	402.0	107.2	838.7
<b>SEd</b>	<b>15.6</b>	<b>17.7</b>	<b>4.76</b>	<b>35.7</b>	<b>12.6</b>	<b>15.1</b>	<b>8.74</b>	<b>29.1</b>
<b>CD (p=0.05)</b>	<b>31.4</b>	<b>35.6</b>	<b>9.60</b>	<b>72.0</b>	<b>25.4</b>	<b>30.5</b>	<b>17.6</b>	<b>58.7</b>

**Table 3. Date of sowing and method of planting on panicle production hill<sup>-1</sup> at harvest stages of rice**

Treatments	Panicles hill <sup>-1</sup>				Panicles m <sup>-2</sup>
	Primary panicle	Secondary panicle	Tertiary panicle	Total	
<b>Main plot : Date of sowing</b>					
D <sub>1</sub>	6.21	5.26	0.18	11.6	402.8
D <sub>2</sub>	5.80	4.98	0.15	10.9	357.9
<b>Mean</b>	6.00	5.12	0.17	11.0	380.4
<b>SEd</b>	<b>0.13</b>	<b>0.09</b>	<b>0.01</b>	<b>0.13</b>	<b>8.06</b>
<b>CD (p=0.05)</b>	<b>NS</b>	<b>0.25</b>	<b>NS</b>	<b>0.56</b>	<b>34.7</b>
<b>Sub plot : Method of planting/sowing with different varieties</b>					
T <sub>1</sub>	6.51	5.63	0.17	12.3	467.0
T <sub>2</sub>	6.20	5.26	0.15	11.4	392.2
T <sub>3</sub>	5.92	4.77	0.11	10.8	323.9
T <sub>4</sub>	6.06	5.31	0.18	11.6	430.5
T <sub>5</sub>	5.98	5.07	0.16	11.2	340.0
T <sub>6</sub>	5.51	4.67	0.15	10.3	298.5
<b>Mean</b>	6.06	5.12	0.17	11.0	380.4
<b>SEd</b>	<b>0.29</b>	<b>0.35</b>	<b>0.01</b>	<b>0.35</b>	<b>19.8</b>
<b>CD (p=0.05)</b>	<b>0.58</b>	<b>0.72</b>	<b>0.02</b>	<b>0.71</b>	<b>39.9</b>

**Table 4. Date of sowing and method of planting with varieties on grain and straw yield**

Treatment	Grain yield (kg ha <sup>-1</sup> )						Straw yield (kg ha <sup>-1</sup> )							
	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	T <sub>6</sub>	Mean	T <sub>1</sub>	T <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	T <sub>5</sub>	T <sub>6</sub>	Mean
D <sub>1</sub>	6291	5323	4537	5458	5021	4183	<b>5136</b>	9680	8571	7125	8967	7954	6477	<b>8129</b>
D <sub>2</sub>	5683	5133	4350	5366	4829	3983	<b>4891</b>	8596	7859	6782	8601	7610	6167	<b>7602</b>
<b>Mean</b>	<b>5987</b>	<b>5228</b>	<b>4443</b>	<b>5413</b>	<b>4925</b>	<b>4083</b>	<b>5013</b>	<b>9869</b>	<b>8946</b>	<b>7684</b>	<b>9515</b>	<b>8513</b>	<b>7053</b>	<b>8597</b>
	<b>D</b>	<b>T</b>	<b>D at T</b>	<b>T at D</b>	<b>D</b>	<b>T</b>	<b>D at T</b>	<b>T at D</b>	<b>D</b>	<b>T</b>	<b>D at T</b>	<b>T at D</b>	<b>D</b>	<b>T</b>
<b>SEd</b>	54.7	173.8	231	198	104.6	274.8	415.9	465.8						
<b>CD (p=0.05)</b>	235.6	350.4	NS	NS	449.8	553.9	NS	NS						



**Fig. 1. Planting methods with different varieties on panicles m<sup>-2</sup> and grain yield (kg ha<sup>-1</sup>)**

At the harvest stage, primary tiller production was more in direct sowing than in unpuddled transplanted rice. A higher number of primary tillers m<sup>-2</sup> (384.9) were recorded from direct sowing of CO 51 and lower number of primary tillers of 296.9 was observed in UPTR of ADT 48. More secondary tillers (458.7) were observed under direct sowing of CO 51 and it was on par with UPTR of CO 51. A lesser secondary tiller m<sup>-2</sup> was recorded from ADT 48 under both methods of planting. With regards to tertiary tillers m<sup>-2</sup>, UPTR of CO 51 produced higher tertiary tillers (125.5) and ADT 48 under direct sowing produced lower tertiary tillers m<sup>-2</sup> (90.6).

Regarding total tillers m<sup>-2</sup>, more number of tillers was recorded in CO 51 rice variety at both methods of Direct sowing (T<sub>1</sub>) and unpuddled transplanting (T<sub>4</sub>). However, comparable tiller production was obtained in MDU 5 rice variety under direct sowing method. But less number of tillers per unit area was observed in ADT 48 under both methods of planting at all the growth stages. The difference in number of tillers m<sup>-2</sup> among the varieties grown under unpuddled condition was also reported by Reddy et al. [6]. More number of tillers per unit area in CO 51 rice might be attributed to the exposure of optimum solar radiation and occurrence of uniform distribution of rainfall during crop growth period, which positively increase the tiller production. The association of increased temperature with low light intensity cause some tillers bud to

dormant due to the non-availability of adequate carbohydrate required for growth [7,8], which restricted the primary, secondary and tertiary production of rice plants. The treatment interaction effect also showed a non significant result at all the stages of observation.

### 3.2 Productive Tillers Hill<sup>-1</sup>

The data on number of productive tillers hill<sup>-1</sup> (primary, secondary, tertiary panicles hill<sup>-1</sup>) at harvest stage is given in Table 3. There was no significant impact on the number of primary and tertiary panicles hill<sup>-1</sup> with respect to sowing dates. Interestingly significant response where observed in secondary panicle production. However, the method of plantings with different rice varieties showed significant variation on number of panicles hill<sup>-1</sup>. Number of primary panicles was more in direct sowing and number of tertiary tillers hill<sup>-1</sup> was more in UPTR method.

Among the treatments, direct sowing of CO 51 (T<sub>1</sub>) produced more primary panicles (6.51) and it was on par with direct sowing of MDU 5 (T<sub>2</sub>) and UPTR of CO 51 (T<sub>4</sub>). The rice variety ADT 48 under direct sowing and UPTR (T<sub>3</sub> and T<sub>6</sub>) recorded 5.92 and 5.51 primary panicles hill<sup>-1</sup>. The same sequence was noticed on secondary panicles hill<sup>-1</sup>. With respect to tertiary panicles, CO 51 under UPTR condition (T<sub>4</sub>) recorded higher value of 0.18 tertiary panicles hill<sup>-1</sup> which was on par with direct sowing of CO 51 (T<sub>1</sub>) and

the lesser tertiary panicles (0.11) was found under UPTR of ADT 48 ( $T_6$ ). These results clearly indicated that direct sowing favoured the production of primary and secondary panicles. Whereas, transplanting favours the production of tertiary panicles because of higher field duration. Moreover, the tertiary panicles remain green and unfilled even the crop attains maturity. Gill et al. [4] also stated similar such research findings.

The number of panicles  $\text{hill}^{-1}$  and per unit area was significantly influenced by date of sowing. Among the sowing dates, the crop sown at July first fortnight ( $D_1$ ) produced maximum number of panicles  $\text{hill}^{-1}$  and per unit area (11.6 and 402.8) than July second fortnight sowing ( $D_2$ ). Regarding method of planting with different varieties, higher panicles  $\text{hill}^{-1}$  (12.3) was recorded under direct sowing of CO 51 ( $T_1$ ) and it was comparable with UPTR of CO 51 ( $T_4$ ) (11.6). ADT 48 under direct sowing and UPTR ( $T_3$  and  $T_6$ ) registered less number of panicles  $\text{hill}^{-1}$  (10.8 and 10.3 respectively). Total number of panicles per unit area was also higher in direct sowing of CO 51 ( $T_1$ ) (467). While, less number of panicles per unit area was obtained in UPTR of ADT 48 ( $T_6$ ) and this was on par with direct sowing of ADT 48 ( $T_3$ ). The interaction effect was not significant with respect to primary, secondary and tertiary panicles  $\text{hill}^{-1}$  and number of panicles  $\text{m}^{-2}$ .

### 3.3 Grain Yield and Straw Yield ( $\text{kg ha}^{-1}$ )

The results on grain yield and straw yield are given in Table 4. The sowing dates as well as method of planting with different varieties brought about significant impact upon the production of grain yield  $\text{kg ha}^{-1}$ . Among the sowing dates, July first fortnight sowing ( $D_1$ ) produced a grain yield of 5136  $\text{kg ha}^{-1}$ , being superior to July second fortnight sowing ( $D_2$ ), which recorded the grain yield of 4891  $\text{kg ha}^{-1}$ . These results were in conformity with the results of Rai and Kushwaha [9] who reported that early sowing crop produced more yield than late sowing, which might be due to optimum period available for growth and development resulted in more storage of photosynthates in the grain in early sown crop. With regards to method of planting with different varieties, direct sowing of CO 51 ( $T_1$ ) produced significantly higher grain yield of 5987  $\text{kg ha}^{-1}$  than other varieties which was followed by UPTR of CO 51 ( $T_4$ ) and direct sowing of MDU 5 ( $T_2$ ) produced a grain yield of 5413 and 5228  $\text{kg ha}^{-1}$ , respectively. The UPTR of ADT 48 ( $T_6$ ) found to

be the lowest yielder (4083  $\text{kg ha}^{-1}$ ) among the treatments.

The results from the Table 4 indicated that, date of sowing and varieties showed a significant effect on this parameter. With respect to sowing dates, the crop sown at July first fortnight ( $D_1$ ) recorded maximum straw yield of 8130  $\text{kg ha}^{-1}$  and this was significantly higher with sowing at second fortnight of July ( $D_2$ ). Among the treatments, direct sowing of CO 51 ( $T_1$ ) produced significantly higher straw yield of 9138  $\text{kg ha}^{-1}$  followed by UPTR of CO 51 ( $T_4$ ) produced 8784  $\text{kg ha}^{-1}$  straw yield it was comparable with direct sowing of MDU 5 (8215  $\text{kg ha}^{-1}$ ). The lower straw yield of 6953 and 6322  $\text{kg ha}^{-1}$  was obtained from ADT 48 under direct sowing and UPTR method ( $T_3$  and  $T_6$ ), respectively. The rate of increase in straw yield proceeded at a higher magnitude in direct sowing of CO 51 rice than the other treatments. Production of more tillers and high accumulation of dry matter would have favoured increased straw yield in CO 51 rice variety than MDU 5 and ADT 48 rice varieties. This is in accordance with the findings of Manjunatha [10]. Direct sown crops produced higher straw yield than UPTR crop and this may be due to more number of tillers per unit area which confirms the findings of Akbar et al. [11] and Praveen et al. [12].

## 4. CONCLUSION

Based on the results, the study concluded that the CO 51 rice proved its supremacy under unpuddled conditions by registering higher growth and yield parameters and yield followed by MDU 5 rice variety. Transplanting under unpuddled condition is found to be comparable with direct sowing conditions and it may be recommended for heavy weed infested area. Regarding to the sowing window, the water released from the reservoir up to second fortnight of July month, direct sowing of short and early duration rice varieties like CO 51 and MDU 5 under unpuddled condition is an ideal for getting higher yield.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Shetty PK, Hedge MR, Mahadevappa M. Innovations in rice production. National

- Institute of Advanced Studies, Bangalore, India; 2013.
2. Natarajan G. Studies on rice based cropping pattern under delayed water release in 2004. (PhD), Tamil Nadu Agricultural University, Coimbatore; 1993.
  3. Belder P, Bouman B, Spiertz J, Peng S, Castaneda A, Visperas R. Crop performance, nitrogen and water use in flooded and aerobic rice. *Plant and Soil*. 2005;273(1-2):167-182.
  4. Gill M, Kumar A, Kumar P. Growth and yield of rice (*Oryza sativa*) cultivars under various methods and times of sowing. *Indian Journal of Agronomy*. 2006;51(2): 123-127.
  5. Gomez KA, Gomez AA. Statistical procedures for agricultural research. 2<sup>nd</sup> Edn. John Wiley and Sons, New York. 2010;680.
  6. Reddy MM, Padmaja B, Veeranna G, Reddy D, Vardhan V. Evaluation of popular kharif rice (*Oryza sativa* L.) varieties under aerobic condition and their response to nitrogen dose. PART I: *Plant Science*. 2012;3(67):37-43.
  7. Yoshida S. Fundamentals of rice crop science. Los Banos, Philippines, IRRI. 1981; 267.
  8. Balajinaik B, Sreenivas G, Raji Reddy D, Leel Rani P. Tillerig behaviour and yield of different duration rice varieties under different dates of sowing under aerobic culture. *International Journal of Current Research*. 2016;8(5):30143–30146.
  9. Rai HK, Kushwaha HS. Effect of planting dates and soil water regimes on growth and yield of upland rice. *Oryza*. 2008;45(1):129-132.
  10. Manjunatha BN, Basavarajappa R, Paramesh V, Anil Kumar S. Impact of age of seedling under different system of rice (*Oryza sativa*) intensification on water used, water use efficiency of economics. Paper presented at the National symposium on Resource management strategies for livelihood Security, Bengaluru, Karnataka, India. 2010;47-52.
  11. Akbar N, Iqbal A, Khan HZ, Hanif MK, Bashir MU. Effect of different sowing dates on the yield and yield components of direct seeded fine rice (*Oryza sativa* L.). *Journal of Plant Breeding and Crop Science*. 2010;2(10): 312-315.
  12. Praveen K, Patel S, Choudhary J, Bhelawe S. Heat unit requirement of different rice varieties under chhattisgarh plain Zones of India. *Journal of Earth Science & Climatic Change*. 2014;5(1):1.

© 2022 Srimathi and Subramanian; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/89853>