



Effect of Different Types of Mulches on Yield of Pumpkin in Salt Affected Soil

Amarendranath Biswas^{a#}, Ajay Kumar Biswas^{b†}
and G. M. Mostafizur Rahman^{c#}

^a Salinity Management and Research Centre, Soil Resource Development Institute, Batiaghata, Khulna, Bangladesh.

^b Soil Resource Development Institute, Divisional Laboratory, Daulatpur, Khulna, Bangladesh.

^c Soil Resource Development Institute, Regional Office, Jashore, Bangladesh.

Authors' contributions

This work was carried out in collaboration among all authors. The author AB came up with the study's plan, ran the stats, drafted the protocol and wrote the first draft of the paper. The authors AKB and GMMR oversaw the literature searches, research analysis and prepared the final paper for submission. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJAHR/2022/v9i4197

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91829>

Original Research Article

Received 17 July 2022
Accepted 21 September 2022
Published 23 September 2022

ABSTRACT

For reducing soil salinity and obtaining a sustainable yield, a low-cost and farmer-friendly method is required for pumpkin, a well-liked vegetable. Accordingly, a field experiment was carried out in Salinity Management and Research Center, Soil Resource Development Institute, Batiaghata, Khulna during Kharif-1 season of 2021 to investigate the effect of different types of mulches on soil salinity and yield of pumpkin. The experiment includes three treatments viz. no mulch (control), straw mulch and mulching paper. The experiment was carried out in Randomized Complete Block Design with four replications. Soil salinity was recorded at 30 days intervals. All mulching materials effectively reduced the salt accumulation in the root zone. After three months of seed sowing, the highest soil salinity (11.32 dS/m) and lowest soil salinity (6.24 dS/m) were found at no mulch (control) and mulching paper treatment respectively in the month of May. Mulching treatments markedly increased the growth and yield attributes of pumpkins. The highest value of three growth parameters i.e., fruit length (26.92 cm), fruit diameter (94.09 cm) and fruit size (2037 cm²) was

[#] Principal Scientific Officer;

[†] Scientific Officer;

*Corresponding author: Email: biswasajay.ku@gmail.com;

found on mulching paper as compared to control. Again, the highest flesh thickness (3.94 cm), fruit weight (4.73 kg) and yield (31.18 t/ha) were recorded at mulching paper treatment whereas the lowest yield (16.68 t/ha) was found at no mulch (control). The results revealed that the use of mulching paper decreased salinity and also increased the yield of pumpkin in saline soil. These findings suggest that the application of mulching paper not only reduces soil salinity but also increases the yield of pumpkins.

Keywords: Salinity; straw mulch; mulching paper; pumpkin; yield; sweet gourd.

1. INTRODUCTION

"In Bangladesh, Pumpkin (*Cucurbita moschata* Duchesne ex Poir.) or sweet gourd is a fairly common and well-liked vegetable. It is grown widely from homesteads to commercial fields and marketed all over the country. Due to its high concentration of vitamin A, spinach is healthy and can help address the lack of vegetables and the nutritional issue. *Cucurbita moschata* is medium-sized and widely grown in the tropics and subtropics, its fruit color is most often dark green, light yellow, or pale orange" [1]. "Pumpkin is a low-calorie vegetable that contains many anti-oxidants such as vitamin-A, vitamin-C and vitamin-E and dietary fiber" [2]. "At present, the total production of vegetables in Bangladesh is 3.73 million tonnes and per capita consumption is currently 62 gm a day against the Food and Agricultural Organization's (FAO) recommendation of 220 gm" [3].

"The foundation of Bangladesh's economy is agriculture which accounts for 12.65% of the country's GDP. Only 7% of the total cropped land in Bangladesh is used for horticulture crops, such as vegetables, while rice accounts for around 75% of the country's agricultural production. The area under vegetable cultivation accounts for only 2.56 percent of the total cropped areas" [4]. "In winter, a good number of vegetables are grown in Bangladesh among them pumpkin or sweet gourd is appreciated by consumers due to their widely usable nature. The Bangladeshi coastal region's agriculture is constantly under threat because of the greater levels of soil and water salinity" [5]. "Both magnitude and levels of soil salinity have been increasing with time being 0.83, 1.02 and 1.06 mha in 1973, 2002 and 2009, respectively" [6]. "In comparison to floodplain agriculture, the average cropping intensity (%) in coastal areas has not increased at the same rate. In the coastal region, between 30 and 50 percent of net cultivated land is left fallow during the Rabi and Kharif-1 seasons. In such regions, intensive irrigation with light saline

surface water exacerbated the issue since the evaporation of the large salt deposits left behind caused secondary salinization and alkalinization" [7].

The salinity rises throughout the dry months, reaching a peak in April-May, and falls during the rainy months, reaching a minimum in July–August [8]. Polythene mulch is utilized extensively worldwide as a crucial farming strategy because of the enormous advantages it provides for increasing crop productivity and water conservation [9]. "The use of polythene mulch may be able to lessen the amount and frequency of irrigation needed for maize production as well as the capillary rise and evaporation losses of water from the soil" [10]. Additionally, it is said that mulching paper is a successful method for encouraging crop emergence because it alters the soil microclimate by raising soil temperature in the winter [10]. "Mulching paper can hold precipitation, decrease water loss and increase the water use efficiency of crop plants. Although some research on using polythene mulch to stop soil erosion and conserve water has been documented" [11], "little is known about how mulch affects saline soil management for potential pumpkin production. It has been shown that the use of mulching paper increased different growth parameters like plant height, the number of leaves per plant, root and shoot fresh and dry weights to a greater extent in maize in salt-affected soil. Additionally, it has been found that the usage of polythene mulch in maize production reduces salinity levels" [12].

Therefore, mulching could be an effective choice for reducing soil salinity and increasing production in coastal areas. Organic mulches can reduce the effect of salt toxicity on plant growth or actively accelerate soil desalinization. Considering the above fact, this experiment is designed to find out the effect of different mulches on soil salinity and yield performance of sweet gourd.

2. MATERIALS AND METHODS

2.1 Study Site

“This experiment was conducted at the Salinity Management and Research Centre (SMRC), Soil Resource Development Institute, Batiaghata, Khulna, Bangladesh during the Kharif-1 season of 2021. Geographically, the study site was at 22°46'01.8" N latitude and 89°24'15.2" E longitude and under AEZ-13. With an average yearly temperature of 79.3 °F and monthly mean temperatures ranging from 54.3 °F in January to 93.7°F in May, the area is among the warmest in Bangladesh” [13].

2.2 Experimental Layout and Crop Management

The experiment was carried out in Randomized Complete Block Design with four replications. Sweet gourd (variety Bengal Sweet-2) was taken as an experimental crop. Three experimental treatments were considered: (a) control (no mulch) (b) straw mulch (c) mulching paper. To achieve good tilth, the experiment plot was prepared by several ploughing and cross ploughing followed by laddering and harrowing with tractor and power tiller. Weeds and other stables were removed carefully from the experimental plot and leveled properly. Manure and basal dose of fertilizer were applied during land preparation. Germinated seeds were sown and necessary intercultural operations were done when necessary. The seeds were sown on 05-02-2021. Data were recorded in accordance with the requirements. Soil salinity was measured by using an EC Meter at 30 days intervals. All the intercultural operations like watering, gap filling, staking, weeding, and plant protection measures were executed carefully.

2.3 Measurement of Growth and Yield Characteristics

Five growth and yield parameters such as fruit length (cm), fruit diameter (cm), fruit size (cm²),

flesh thickness (cm) and fruit weight (kg) were taken into consideration to analyze the effect of mulches on yield of sweet gourd. Total yield (t/ha) was calculated by measuring the total fruit weight of a plot.

2.4 Statistical Analysis

The collected data were tabulated and statistically analyzed using Statistix 10 software. The treatment means were separated statistically at a 5 % level of significance using Duncan’s Multiple Range Test (DMRT).

3. RESULTS AND DISCUSSION

3.1 Effects of Different Mulches on Soil Salinity

“Control of root zone salinity in saline fields is considered beneficial to seed emergence and stand establishment” [14]. “Previous studies have shown that mulching with different materials is a promising technique for salinity control in agriculture” [15]. An upward trend of soil salinity at all treatments was observed from February to May (Table 1). In the month of February, the lowest soil salinity was found at mulching paper (4.80 dS/m) while the highest soil salinity was observed at control (5.62 dS/m) where no mulch was used. Since soil salinity increases gradually from the month of February to May, the lowest salinity was found in the month of February (4.80 dS/m) as compared to the highest in the month of May (11.32 dS/m). In the month of May, the lowest soil salinity was found at mulching paper (6.24 dS/m) in comparison with the highest soil salinity found in control (11.32 dS/m) (Table 1). “Thus, it revealed that soil salinity can be reduced by using different mulches in which mulching paper has a significant effect on reducing soil salinity. These results are emphasized by some authors who reported that all mulching materials effectively reduced salt accumulation in the root zone” [16].

Table 1. Effect of mulches on soil salinity of experimental plot during cultivation period

Treatment	Month-wise soil salinity (EC: dS/m)			
	Feb	Mar	April	May
Control	5.62	7.65	9.59	11.32
Straw Mulch	5.24	6.25	8.62	9.25
Mulching Paper	4.80	5.32	6.85	6.24

Source: Soil Salinity Test Results

3.2 Effects of Different Mulches on Growth Characteristics of Pumpkin cv. Bengal Sweet-2

“Fruit length, fruit diameter and fruit size were statistically analyzed as shown in Table 2. They showed a significant variation in relation to different mulch materials. The highest fruit length (26.92 cm) was found at mulching paper while the lowest (16.04 cm) was found at control where no mulch was applied. The highest fruit diameter (94.09 cm) and fruit size (2037 cm²) were observed at mulching paper in comparison with the control where the lowest fruit diameter (66.03 cm) and fruit size (1638 cm²) were found (Table 2). Sageer Khan et al (2015) found that fruit length (cm), fruit weight (g), fruit diameter (cm), no. of fruits per plant and fruit yield (213.47q/ha) of sponge gourd were significantly superior to black polyethylene mulch while, plants without mulch (control) resulted in poor growth and yield” [17]. “These results are in conformity with Islam et al (2021) who reported that yield characteristics like diameter of fruit, weight of fruit and fruits per vine showed significant results with black polyethylene mulch in case of different high-value vegetables in Bangladesh” [18]. “A similar trend was also found in the findings of Yaseen et al (2014) who reported that leaf area (LAI) and plant height were significantly affected by the mulching treatments” [19].

3.3 Effects of Different Mulches on Yield Attributes of Pumpkin cv. Bengal Sweet-2

Flesh thickness, fruit weight and yield of sweet gourd grown under different mulching treatments are presented in Table 3. Statistical analysis was carried out on yield and yield attributes which revealed that these are significantly varied due to different mulches. Maximum flesh thickness (3.94 cm) was recorded in mulching paper which was statistically similar to straw mulch (3.79 cm). While minimum flesh thickness (2.93 cm) was observed in control. Again, the highest fruit weight (4.73 kg) was recorded in mulching paper followed by straw mulch (3.83 kg) whereas the lowest fruit weight (2.15 kg) was found in control treatment (Table 3). The researchers indicated that plants under polyethylene mulch produce larger fruit and have higher fruit yield per plant because of the better plant growth that is due to a favorable hydro-thermal regime of soil and a completely weed-free environment. The data depicted that the highest yield (31.18 t/ha) was found at mulching paper while the lowest (16.68 t/ha) was found at control where no mulch was applied. By encouraging changes in the plant's microenvironment, plastic mulch enhances plant growth and vigor as well as production and yield.

Table 1. Effect of mulches on yield and yield contributing characters of Pumpkin

Treatment	Fruit Length (cm)	Fruit diameter (cm)	Fruit size (cm ²)
Control	16.04c	66.03c	1638c
Straw Mulch	23.82b	82.98b	1816b
Mulching Paper	26.92a	94.09a	2037a
Level of sig.	**	**	**
CV (%)	6.80	4.19	3.33

Level of significance ** = 1%; * = 5%.

Table 3. Effect of mulches on yield and yield contributing characters of Pumpkin

Treatment	Flesh thickness (cm)	Fruit weight (kg)	Yield (t/ha)
Control	2.93b	2.15c	16.68c
Straw Mulch	3.79a	3.83b	20.62b
Mulching Paper	3.94a	4.73a	31.18a
Level of sig.	*	**	**
CV (%)	4.93	7.02	9.32

Level of significance ** = 1%; * = 5%. Means having similar letter(s) are statistically similar and those having different letter(s) differ significantly at 5% level

A similar result was reported by Mahadeen (2014) who observed that polythene mulch resulted in higher yield in two summer vegetables compared to straw mulch and no mulch [20]. "This result may be due to the improvement of soil physical properties as well as increasing soil water holding capacity which gave rise to good aeration and drainage that encourage better root growth and nutrient absorption" [20]. "This result is in the line with those recorded by Yonggan et al (2016) who reported that the combined application of straw layer burial and plastic mulch performed better than either the individual use of straw layer or plastic mulch for controlling salt accumulation in the topsoil layer and increasing sunflower yield" [21]. Akhter et al (2018) also reported that "black polythene mulch and phosphorus at 90 kg P₂O₅/ha would be suitable for optimum vegetative growth and yield of squash compared to other treatments" [22]. Ibraheem et al (2019) also found that soil treated with black plastic mulch exhibited a substantial increase in leaf area, fruits number, fruits weight, fruits length, and overall yield of summer squash in comparison to plants planted in non-mulching soil [23]. "The findings of Farah et al (2021) also showed a similar pattern. They found that plastic mulch dramatically improved stomatal conductance, photosynthesis, transpiration, leaf chlorophyll fluorescence, and leaf chlorophyll levels, which resulted in a 34% increase in squash production and a 46% increase in irrigation water use efficiency under partial root drying plants" [24].

4. CONCLUSION

It may be concluded from the experiment's findings that using mulching materials led to a noticeable decrease in the accumulation of salts. Maximum soil salinity was reduced by using mulching paper in comparison with the control treatment where no mulch was applied. The reduction of soil salinity followed the following order of treatment: mulching paper > straw mulch > control. Different growth and yield attributes were significantly impacted due to different mulching treatments. Results implied that mulching paper treatment gave the highest yield (31.18 t/ha) followed by straw mulch treatment yielded 20.62 t/ha. Whereas, the lowest yield (16.68 t/ha) was recorded in control treatment. Between two types of mulch mulching paper can be used at the farmer's level to reduce soil salinity and increase the yield of sweet gourd. However, further research is still needed to work

out a cost-effective technology to reduce soil salinity and increase the yield of sweet gourds.

ACKNOWLEDGEMENTS

For the successful completion of the study, the authors would like to express their special gratitude to the Salinity Management and Research Centre (SMRC), Soil Resource Development Institute (SRDI), Batiaghata, Khulna, Bangladesh and the research team members.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Kiramana JK, Isutsa DK. First detailed morphological characterization of qualitative traits of extensive naturalized pumpkin germplasm in Kenya. *International Journal of Development and Sustainability*. 2017;6(7):500-525.
2. Yadav M, Jain S, Tomar R, Prasad GB, Yadav H. Medicinal and biological potential of pumpkin: An Updated Review. *Nutrition Research Reviews*. 2010;23(2):184–190.
3. Anonymous. Per capita deficiency of vegetables 158gm a day. *The Financial Express*; 2022. Available: <https://rb.gy/t7rwbh>.
4. BBS. Yearbook of Agricultural Statistics of Bangladesh, (33rd Series). Bangladesh Bureau of Statistics, Ministry of Planning, Government of the People's Republic of Bangladesh; 2021. Available: <https://tinyurl.com/49c4dhda>.
5. MoA and FAO. Master Plan for Agricultural Development in the Southern Region of Bangladesh. Ministry of Agriculture; 2013;37-40. Available: <https://www.fao.org/3/au752e/au752e.pdf>.
6. Fertilizer Recommendation Guide (FRG). Bangladesh Agricultural Research Council (BARC), Farmgate, Dhaka. 2012;66.
7. Roychoudhury A, Chakraborty M. Biochemical and molecular basis of varietal difference in plant salt tolerance. *Annual Review & Research in Biology*. 2013;3(4):422-454.
8. Haque MA, Jahiruddin M, Hoque MA, Rahman MZ, Clarke D. Temporal

- variability of soil and water salinity and its effect on crop at Kalapara upazila. *Journal of Environmental Science & Natural Resources*. 2014;7(2):111-114.
9. Sun SJ, Fan YM, Xu ZH, Zhang XD, Chi DC. Effects of planting density on soil moisture and corn yield under plastic film mulching in a rain-fed region of northeast China. *Chinese Journal of Ecology*. Chinese.2014;33:2650– 2655.
 10. Bu L, Liu J, Zhu L, Luo S, Chen X, Li SQ. The effects of mulching on maize growth, yield and water use in a semi-arid region. *Agriculture Water Management*. 2013; 123:71–78.
 11. Zhang G, Zhang X, Hu X. Runoff and soil erosion as affected by plastic mulch patterns in vegetable field at Dianchilake's catchment, China. *Agricultural Water Management*. 2013;122: 20-27.
 12. Sikder MU, Haque MA, Jodder R, Kumar T, Mondal D. Polythene Mulch and Irrigation for Mitigation of Salinity Effects on Maize (*Zea mays* L.). *The Agriculturists*. 2016;14(2):1-13.
 13. Anonymous. Bangladesh agro-meteorological information portal. Department of Agricultural Extension; 2022. Available:<https://www.bamis.gov.bd/en/page/aezs-maps/>
 14. Dong H, Li W, Xin C, Tang W, Zhang D. Late planting of short-season cotton in saline fields of the Yellow River Delta. *Crop Sci*. 2010;50:292–300.
 15. Pang H, Li Y, Yang J, Liang Y. Effect of brackish water irrigation and straw mulching soil salinity and crop yields under monsoonal climatic conditions. *Agric. Water Manag.* 2010;97:1971-1977.
 16. Taia A. Abd El-Mageed, Wael M. Semida, Mohamed H. Abd El-Wahed. Effect of mulching on plant water status, soil salinity and yield of squash under summer-fall deficit irrigation in salt affected soil. *Agricultural Water Management*. 2016; 173(C):1-12. Available:<https://doi.org/10.1016/j.agwat.2016.04.025>.
 17. Khan S, Pal M, Kumar V. Influence of different mulches on growth and yield of sponge gourd (*Luffa cylindrica* L.). *Plant Archives*. 2015;15(1):393-395.
 18. Islam F, Quamruzzaman A, Mallick SR. Effect of different mulch paper on growth and yield of different high value vegetables in Bangladesh. *Agricultural Sciences*. 2021;12:237-246. Available:<https://doi.org/10.4236/as.2021.123015>
 19. Yaseen R, Shafi J, Ahmad W, Rana MS, Salim M, Qaisrani SA. Effect of Deficit Irrigation and Mulch on Soil Physical Properties, Growth and Yield of Maize. *Environment and Ecology Research*. 2014;2(3):122-137. DOI: 10.13189/eer.2014.020302.
 20. Mahadeen AY. Effect of polyethylene black plastic mulch on growth and yield of two summer vegetable crops under rain-feed conditions under semi-arid region conditions. *American Journal of Agricultural Biological Sciences*. 2014; 9(2):202-207.
 21. Yonggan Z, Yuyi L, Jing W, Huancheng P, Yan L. Buried straw layer plus plastic mulching reduces soil salinity and increases sunflower yield in saline soils. *Soil and Tillage Research*. 2016;155: 363-370. DOI:10.1016/j.still.2015.08.019.
 22. Akhter F, Mostarin T, Islam MN, Akhter S. Effect of Mulches and Phosphorus on Growth and Yield of Squash (*Cucurbita pepo*).*The Agriculturists*. 2018; 16:25-34. DOI: 10.3329/agric.v16i02.40340.
 23. Ibraheem FF, Allel WB, Hussien JM. Effect of soil mulching, organic and inorganic fertilizer on growth and yield of summer squash. *International Journal of Agricultural Statical Sciencei*. 2019;15(2): 677-85.
 24. Farah AH, Al-Ghobari HM, Zin El-Abidin TK, Alrasasimah MS, El-Shafei AA. Impact of partial root drying and soil mulching on squash yield and water use efficiency in arid. *Agronomy*. 2021;11(4):706.

© 2022 Biswas et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/91829>