



Efficacy of Bio-Agents in Managing Root Rot of Okra Caused by *Rhizoctonia solani*

Ranjana Meena ^{a*}, R. P. Ghasolia ^{a++}, Shailesh Godika ^{a++},
Raja Ram Bunker ^b, Pinki Devi Yadav ^c, Suman Chopra ^a
and Mateti Gayithri ^d

^a Department of Plant Pathology, SKN College of Agriculture, Jobner-303328 (Jaipur-Rajasthan), India.

^b ICAR-KVK, Hailakandi, ICAR Research Complex for NEH Region, Assam- 78815, India.

^c Rajasthan Agricultural Research Institute, Durgapura, SKNAU, Jobner (Jaipur-Rajasthan), India.

^d Assam Agricultural University, Assam- 785013, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/jeai/2024/v46i113029>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here:

<https://www.sdiarticle5.com/review-history/125734>

Original Research Article

Received: 20/08/2024

Accepted: 22/10/2024

Published: 06/11/2024

ABSTRACT

Okra or lady's finger [*Abelmoschus esculentus* (L.) Moench] is one of the most important summer vegetables of Rajasthan as well as India. Okra is attacked by several diseases caused by fungi and important one is root rot caused by *Rhizoctonia solani* Kuhn, which is an important constraint to the crop and causes significant economic losses. During present study, two bio-agents were evaluated

⁺⁺ Professor;

*Corresponding author: E-mail: ranjanameena985@gmail.com;

Cite as: Meena, Ranjana, R. P. Ghasolia, Shailesh Godika, Raja Ram Bunker, Pinki Devi Yadav, Suman Chopra, and Mateti Gayithri. 2024. "Efficacy of Bio-Agents in Managing Root Rot of Okra Caused by *Rhizoctonia Solani*". *Journal of Experimental Agriculture International* 46 (11):50-56. <https://doi.org/10.9734/jeai/2024/v46i113029>.

alone and in combination *Trichoderma harzianum* through seed (4 g/kg) and soil (4 kg/ha) applications for two consecutive years under artificial inoculation conditions in the field. Among these, seed + soil application of *Trichoderma harzianum* was found most effective in reducing disease incidence (48.76%) and in increasing pod yield (13.46%) followed by seed treatment with *T. harzianum* + soil application of *Pseudomonas fluorescens* (46.41% and 12.84%, respectively). Looking to the increasing demand of organically produces in the market, root rot disease can be managed effectively with these eco-friendly approaches for supplying organically produced vegetables to fulfil the ever-increasing demand among health-conscious people.

Keywords: Okra; root rot; *rhizoctonia solani*; bio-control agents; disease control.

1. INTRODUCTION

“Okra [*Abelmoschus esculentus* (L.) Moench] is an important vegetable valued for its edible green pods. The geographical origin of okra is disputed, with supporters of South Asian, Ethiopian and West African origins. The plant is cultivated in tropical, subtropical and warm temperate regions around the world” (Anonymous, 2006). “It considers as good source of proteins, carbohydrates, vitamins, calcium, enzymes and total minerals, as well as it has a therapeutic effects in the treatment of ulcer and relief of hemorrhoids, and as a substitute for blood plasma. Additionally is useful in the treatment of urinary and reproductive system disorders” (Singh et al., 2014).

“Furthermore, it can be beneficial from its fresh leaves, buds, flowers, horns, stems and seeds, and it is an immature fruit as vegetables. Its seeds are a source of oil, where the oil concentration range from 20% to 40% and contains tryptophan acid up to 47.4%, in the form of unsaturated fatty acid that necessary for human feed. Moreover, Okra helps to reduce cholesterol in blood, the other part of the okra is an insoluble fiber, which helps to maintain intestinal health, and it is rich in phenolic compounds with biological properties such as quaternary and flavonol derivatives, catechin oligomers and hydroxycinnamic derivatives” (Gemedie et al., 2015).

“Okra crop is attacked severely from the vagary of diseases caused by fungi, bacteria, viruses and nematodes in the field, which not only reduces the potency of seed, but also degrades the health beneficial and nutritional quality components of the fruits. The important diseases of okra are root rot (*Rhizoctonia solani* Kuhn), powdery mildew (*Oidium* spp.), Fusarium wilt (*Fusarium oxysporum*), charcoal rot (*Macrophomina phaseolina*), Cercospora leaf spot (*Cercospora abelmoschi*), damping off

(*Pythium* spp.), root knot (*Meloidogyne* sp.) and yellow vein mosaic (*Bhindi Yellow Vein Mosaic Virus*)” (Anonymous, 2003). Among the fungal diseases, root rot is one of the important diseases caused by *Rhizoctonia solani*.

“Amongst these diseases, root rot caused by *Rhizoctonia solani*, is an important constraint to the crop and causes significant losses. The pathogen mainly attacks the root and underground parts, but it is also capable of infecting the other plant parts like the green foliage parts, the seeds and the hypocotyls” (Acharya et al., 2014). “Among the initial symptoms of the disease, yellowing of leaves is a first symptom which in next two or three days, leaves droop and wither off. Infected plants may wilt within a week after the appearance of first symptom. When stem is examined closely, dark lesions can be observed on the bark near ground level. The roots of infected plants are poorly developed; finer roots are either not formed or rotted. Plants show stunted growth and can easily be pulled out. If the plants are pulled from soil, the basal stem along with main root, may show symptoms of rotting. The tissues are weakened and break off easily in advanced cases and sclerotial bodies can be seen scattered on the affected roots. The fungus is mainly a soil dweller and spreads from plant to plant through irrigation water and implements and cultural operations. The sclerotia and pycniospores may also become air borne and cause further spread of the pathogen (Rangaswami and Mahadevan, 2008). Crop losses by root rot of okra (*Rhizoctonia solani*) is ranged from negligible to 50-60 per cent depending on the extent of severity and different stages of crop” (Safiuddin et al., 2014).

“Exclusive dependency on fungicides and their indiscriminate use for managing diseases of various crops has resulted in residue, environmental hazards and development of resistance in pathogens. Due to increasing

consciousness in consumers about organic produce, particularly medicinal products, efforts are being diverted to manage the diseases through eco-friendly approaches as a tool for integrated disease management because they do not cause bio-accumulation in eatables and environmental pollutions. Involvement of bio-agents in disease management system is also an important segment of disease control, looking to the hazards caused by toxic chemicals or in a situation where pathogens develop resistance to fungicides" [Yadav and Ghasolia, 2022]. In lieu of this, the present investigation was undertaken to evaluate different bio-agents alone and in combination in field for two consecutive years for managing the disease and increasing yield recovery for the profit of growers and consumers.

2. MATERIALS AND METHODS

The field experiment was conducted at Instructional Farm, S.K.N. College of Agriculture, Jobner, Jaipur (Rajasthan) for two consecutive years. Jobner is situated at

DMS – Latitude - 26°58'21.25" N
Longitude - 75°23'15.07" E

DD - Latitude - 26.97257°
Longitude - 75.38752°

The region falls under semi-arid eastern plain (Agro Climatic Zone- III A) of Rajasthan.

The two most promising fungal and bacterial bio-agents were evaluated through seed and soil applications for two consecutive years during *Zaid* season 2022 and 2023 either alone or in combinations in randomized block design (RBD) in 1.8 m x 2.25 m plots with three replications. Commercial formulation of bio-agents (*Trichoderma harzianum* and *Pseudomonas fluorescens*) was applied through seed treatment and soil methods. Before applying bio-agents on the surface of the seeds, seeds were moistened with five per cent gum solution applied at 10 ml/kg of seeds. The bio-agents were used as seed treatment @ 4 g/kg seeds where as in soil application @ 4 kg /ha. The crop was sown in first week of Feb. Inoculum multiplied on sorghum grains was added @ 20 g/m row length at the time of sowing. Observations on disease incidence (75 DAS) and pod yield (up to harvest) were recorded.

3. RESULTS AND DISCUSSION

Two most promising bio-agents were evaluated through seed and soil applications for two

consecutive years during *Zaid* 2022 and 2023 either alone or in combinations. These bio-agents were caused significant reduction in root rot incidence and increased pod yield of okra in comparison to untreated check (Table 1 and Fig. 1). Among these, seed + soil application of *Trichoderma harzianum* was found most effective in reducing disease incidence (48.76%) and in increasing pod yield (13.46%) followed by seed treatment with *T. harzianum* + soil application of *Pseudomonas fluorescens* (46.41% and 12.84%), seed treatment with *P. fluorescens* + soil application of *T. harzianum* (45.32% and 10.45%, respectively). The soil application of *P. fluorescens* was found least effective in reducing the disease incidence (18.83%) and in increasing minimum pod yield (3.12%) over control.

Incremental cost benefit ratio (ICBR) was calculated to interpretate the economics of these bio-agents. The data indicated that the highest ICBR was obtained by the seed application of *Trichoderma harzianum* (1:53.52) followed by seed treatment with *P. fluorescens* (1:48.23), seed + soil treatment with *T. harzianum* (1:22.73), *T. harzianum* (seed treatment) + *P. fluorescens* (soil application) (1:21.67), *P. fluorescens* (seed treatment) + *T. harzianum* (soil application) (1:17.64), *P. fluorescens* (seed treatment) + *P. fluorescens* (soil application) (1:16.34) and *T. harzianum* (soil application) (1:7.06) whereas, lowest ICBR was observed with soil application of *P. fluorescens* (1:5.31).

"The significance of chemicals and bio-control agents cannot be denied in plant disease management and especially bio-agents that have been proved to be highly effective in managing soil borne fungal plant pathogens. Our findings are in accordance with the results" of Melo and Faull (Melo and Faull 2000) and Rangeshwaran et al. (2001). Melo and Faull (2000) "screened 14 *Trichoderma* strains for the control of *Rhizoctonia solani* and among them three *T. koningii* strains produced toxic metabolites with strong activity against *Rhizoctonia solani*, inhibiting mycelial growth by 79 per cent. *T. harzianum* and *T. koningii* also reduced the viability of sclerotia of *Rhizoctonia solani* by 81.8 per cent and 53 per cent, respectively. Rangeshwaran et al. (2001) studied the impact of seed treatment with two bacterial antagonists viz., *Pseudomonas putida* and *Pseudomonas fluorescens* to control the root rot of chickpea caused by *Rhizoctonia* sp. and found that seed treatment with both the antagonists significantly reduced root rot of chickpea. Among these

Table 1. Effectiveness of bio-control agents in managing root rot of okra under field conditions

T. No.	Bio agents	PDI		Pooled	Disease reduction (%) over control	Yield (g/ha)		Pooled	Per cent increase in yield over check	Cost of treatment + Labour Charges (ha)	Gross return (ha)	ICBR ratio
		2022	2023			2022	2023					
T1	<i>T. harzianum</i> (seed treatment)	43.12 (41.05)	40.45 (39.49)	41.79 (40.27)	26.89	54.26	52.02	53.14	7.36	272	212560	1:53.52
T2	<i>P. fluorescens</i> (seed treatment)	44.34 (41.75)	41.33 (40.01)	42.84 (40.88)	25.05	53.68	51.87	52.78	6.63	272	211120	1:48.23
T3	<i>T. harzianum</i> (soil application)	45.09 (42.18)	42.33 (40.59)	43.71 (41.39)	23.52	52.84	50.25	51.55	4.14	1160	206200	1:7.06
T4	<i>P. fluorescens</i> (soil application)	47.23 (43.41)	45.55 (42.45)	46.39 (42.93)	18.83	52.05	50.03	51.04	3.12	1160	204160	1:5.31
T5	<i>T. harzianum</i> (seed treatment) + <i>T. harzianum</i> (soil application)	30.44 (33.49)	28.13 (32.03)	29.29 (32.77)	48.76	57.18	55.13	56.16	13.46	1172	224640	1:22.73
T6	<i>P. fluorescens</i> (seed treatment) + <i>Pseudomonas fluorescens</i> (soil application)	35.23 (36.41)	30.33 (33.42)	32.78 (34.93)	42.65	55.46	53.12	54.29	9.69	1172	217160	1:16.34
T7	<i>T. harzianum</i> (seed treatment) + <i>P. fluorescens</i> (soil application)	32.43 (34.71)	28.83 (32.48)	30.63 (33.60)	46.41	57.29	54.41	55.85	12.84	1172	223400	1:21.67
T8	<i>P. fluorescens</i> (seed treatment) + <i>T. harzianum</i> (soil application)	33.35 (35.27)	29.16 (32.68)	31.26 (33.99)	45.32	56.09	53.24	54.67	10.45	1172	218680	1:17.64
T9	Control	59.16 (50.28)	55.15 (47.96)	57.16 (49.12)	0.00	50.61	48.38	49.50	0.00	-	198000	-
	SEm±	1.50	1.41	0.89		2.88	2.69	1.70				
	CD (P=0.05)	4.51	4.24	2.68		8.67	8.10	5.14				
	CV (%)	6.51	6.43	4.75		9.16	8.94	6.64				

*Average of three replications. Figures in parentheses are angular transformed values




Fig. 1. Efficacy of bio-agents on root rot of okra under field conditions

antagonists, *P. fluorescens* was found to be highly effective in reducing (6.4%) incidence as compared to *P. putida*". Madhi et al. (2020) "who conducted study on root rot and damping-off diseases of okra and control them biologically using *Pseudomonas fluorescens* bacteria. The results showed the isolation of several pathogenic fungal, as well as fungi is also accompanying them, where the *Rhizoctonia solani* recorded the highest percentage of appearance and frequency which was (14.6, 16%) respectively. Singh et al. (2020) concluded that microbial volatile organic compounds (mVOCs) have great potential in plant ecophysiology, yet the role of below ground VOCs in plant stress management remains largely obscure. Analysis of biocontrol producing VOCs into the soil allow detailed insight into their interaction with soil borne pathogens for plant disease management. A root interaction trial was set up to evaluate the effects of VOCs released from *Trichoderma viride* BHU-V2 on soil-inhabiting fungal pathogen and okra plant growth". Ahmed (2023) "stated that management of plant root diseases by the application of seaweed, particularly fluorescent *Pseudomonas*, is capturing the interest of plant scientists. In this study, the effect of seaweed soil amendment alone or mixed with endophytic fluorescent *Pseudomonas* (EFP-47) in reducing the root infecting fungi of okra was evaluated in pots and field plot experiments. It is suggested that endophytic fluorescent *Pseudomonas* and seaweed could be used for the management of root diseases of okra".

4. CONCLUSION

Conclusively, seed + soil application of *Trichoderma harzianum* was found most effective in reducing disease incidence (48.76%) and in increasing pod yield (13.46%) followed by seed treatment with *T. harzianum* + soil application of *Pseudomonas fluorescens* (46.41% and 12.84%, respectively), seed treatment with *P. fluorescens* + soil application of *T. harzianum* (45.32% and 10.45%, respectively). The soil application of *P. fluorescens* was found least effective bio-agents in reducing the disease incidence (18.83%) and in increasing minimum pod yield (3.12%) over control.

DISCLAIMER (ARTIFICIAL INTELLIGENCE)

Author(s) hereby declare that NO generative AI technologies such as Large Language Models

(ChatGPT, COPILOT, etc) and text-to-image generators have been used during writing or editing of this manuscript.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Anonymous. (2006). *Okra lost crops of Africa: Volume II: Vegetable 2*. National Academies Press. ISBN 978-0-309-10333.
- Singh, P. V., Chauhan, B. K., Tiwari, S. S., Chauhan, S., Simon, S. B., & Abidi, A. B. (2014). An overview on okra (*Abelmoschus esculentus*) and its importance as a nutritive vegetable in the world. *International Journal of Pharmacy and Biological Sciences*, 76(4), 227-233.
- Gemede, H. F. N., Ratta, G. D., Haki, A. Z., Woldegiorgis, M., & Beyene, F. (2015). Nutritional quality and health benefits of okra (*Abelmoschus esculentus*): A review. *International Journal of Nutrition and Food Sciences*, 4(2), 208-215.
- Anonymous. (2003). *Indian Agriculture* (pp. 271-272). Vikas Singhal for Indian Economies Data Research Centre, Maya Puri, New Delhi.
- Acharya, K., Chakraborty, N., Chatterjee, S., & Basu, S. K. (2014). Fungal diseases of fenugreek. *American Journal of Social Issues and Humanities*, 176.
- Rangaswami, G., & Mahadevan, A. (2008). *Diseases of crop plants in India* (4th ed.). Prentice-Hall of India Private Limited.
- Safiuddin, S. A., Tiyagi, R. R., & Mahmood, I. (2014). Biological control of disease complex involving *Meloidogyne incognita* and *Rhizoctonia solani* on growth of okra through microbial inoculants. *Journal of Microbiology and Biotechnology Research*, 4(5), 46-51.
- Yadav, S. L., & Ghasolia, R. P. (2022). Management of root rot of fenugreek caused by *Rhizoctonia solani* using plant extracts and bio-agents. *Legume Research-An International Journal*, 45(3), 391-395.
- Melo, I., S. De, & Faull, J. L. (2000). Parasitism of *Rhizoctonia solani* by strains of *Trichoderma* spp. *Scientia Agricola*, 57(1), 55-59.
- Rangeshwaran, R., Prasad, R. D., & Anuroop, C. P. (2001). Field evaluation of two bacterial

- antagonists *Pseudomonas putida* against wilt and root rot of chickpea. *Journal of Biological Control*, 15, 165-170.
- Madhi, Q. H., Dagher, G. M., & Jumaah, A. M. (2020). Investigation of the causes of root rot and damping-off diseases of okra and testing the effectiveness of *Pseudomonas fluorescens* in disease control. *Plant Archives*, 20(1), 1361-1366.
- Singh, J., Singh, P., Vaishnav, A., Ray, S., Rajput, R. S., Singh, S. M., & Singh, H. M. (2020). Belowground fungal volatiles perception in okra (*Abelmoschus esculentus*) facilitates plant growth under biotic stress. *Microbiological Research*, 246, 126-721.
- Ahmed, L., Zehra, R., Rahman, S. U., Moin, S., Enamullah, S. M., & Haque, S. E. (2023). Enhancement of defensive capability of okra (*Abelmoschus esculentus* L., Moench) against root rot disease by the endophytic fluorescent *Pseudomonas* and seaweed soil amendment. *Pakistan Journal of Botany*, 55(3), 1187-1196

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of the publisher and/or the editor(s). This publisher and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/125734>